

Impact Of Sugars On Curd Bacteria (*Lactobacillus*)

Monalisa Kulshrestha*, Ajaya Eesha and Meera Gupta

*Department of Botany,
S.S. Jain Subodh Autonomous PG College, Jaipur, Rajasthan, India
Corresponding author: monalisakulshrestha@gmail.com

Abstract

The present investigation is based on the effect of different types of sugars on the growth of curd bacteria in vitro. It was observed that upon the utilization of sucrose, dextrose, fructose and lactose; most potential sugar type was lactose for the growth of curd bacteria. This was followed by fructose.

Key words: Curd, Bacteria, Nutrient agar, Growth, Sugars, Lactobacillus, Culture, Probiotic

INTRODUCTION

The present study is based on the increment of curd bacterial culture by using different types of sugars-sucrose, lactose, dextrose and fructose. It will be observed which type of sugar best causes increment of curd bacteria and this can be used in large scale for curd production in larger amount. Based on the above criteria, the aims and objectives of the present research are: Obtaining curd bacterial culture using simple nutrient agar (NA) media, addition of sucrose, dextrose, fructose, lactose in curd and make comparative account of increment in curd bacteria in presence of different types of sugars²³.

MATERIALS AND METHODS

Preparation of Nutrient Agar (NA) media

28 gm of NA powder was added to 1000 ml of boiling water and boiled, stirred and autoclaved. This was utilized for the experimentation.

5 samples were prepared:

- a. Normal curd sample
- b. 1ml curd + 0.1 gm sucrose
- c. 1 ml curd + 0.1 gm fructose

- d. 1 ml curd + 0.1 gm lactose
- e. 1 ml curd + 0.1 gm dextrose

Pouring of NA media was done in laminar air flow cabinet after proper sterilization. Solidification was done after pouring of media in petriplates. The 5 samples were added to petriplates with NA media by spread plate method.

- a. Inoculation of 1 ml curd sample in NA by spread plate method
- b. Inoculation of 1ml curd + 0.1 gm sucrose in NA by spread plate method
- c. Inoculation of 1ml curd + 0.1 gm fructose in NA by spread plate method
- d. Inoculation of 1ml curd + 0.1 gm dextrose in NA by spread plate method
- e. Inoculation of 1ml curd + 0.1 gm lactose in NA by spread plate method

The method of spread plating was carried out as follows:

Spread plate technique:

Pipette the required amount of bacteria on the surface of the Petri plate containing media. Spread the curd inoculum over the surface of the agar medium using a spreader L shaped. This spreader is rotated in Petri plate on all sides. Incubate the plate inverted at 37 C. Lawn of bacteria can be seen in Petri plates. The different inoculums used in different Petri plates can be weighed.

Weight of simple NA media containing Petri plate taken was 40.42 gm

Weight of NA media with curd bacteria as inoculums = 41.40 gm

Weight of NA media with curd bacteria containing sucrose as inoculum = 46.50 gm

Weight of NA media with curd bacteria containing fructose as inoculums = 43.70 gm

Weight of NA media with curd bacteria containing dextrose as inoculums = 42.59 gm

Weight of NA media with curd bacteria containing lactose as inoculums = 50.44 gm

Weight of bacteria was calculated = wt of (NA+inoculums) petriplate – wt of NA petriplate

Weight of simple curd cultured in NA = 41.40-40.42

Weight of curd in presence of sucrose = 46.50-40.42

Weight of curd in presence of fructose = 43.70-40.42

Weight of curd in presence of dextrose = 42.59-40.42

Weight of curd in presence of lactose = 50.44-40.42

RESULTS AND DISCUSSION

Lactobacillus, is a genus of Gram-positive, facultative anaerobic or microaerophilic, rod-shaped bacteria. They are a major part of the lactic acid bacteria group²⁵. In humans they are part of the vaginal microbiota^{1,2}. Many species in this genus have had their genomes sequenced^{3,4}.

Lactobacillus is a Gram-positive (it retains crystal violet dye), facultative anaerobe (it can produce energy through glycolysis and fermentation when oxygen is not present).

Lactobacillus is a member of the lactic acid bacteria group (its members convert lactose and other sugars to lactic acid)^{5,6,7}.


Lactobacillus acidophilus (New Latin 'acid-loving milk-bacillus') is a species of gram positive bacteria in the genus *Lactobacillus*. *L. acidophilus* is a homofermentative, microaerophilic species, fermenting sugars into lactic acid, and grows readily at rather low pH values (below pH 5.0) and has an optimum growth temperature of around 37 °C (99 °F). *L. acidophilus* occurs naturally in the human and animal gastrointestinal tract and mouth²⁴. Some strains of *L. acidophilus* may be considered to have probiotic characteristics²². These strains are commercially used in many dairy products, sometimes together with *Streptococcus thermophilus* and *Lactobacillus delbrueckii* subsp. *bulgaricus* in the production of acidophilus-type yogurt. Its genome has been sequenced^{8,9,10}.

***Lactobacillus acidophilus* cell structure**

The structure of *L. acidophilus* has many characteristics. This bacterium is a single-celled prokaryotic microorganism that lacks a distinct nucleus. Although, there are many other parts to the cell of this bacterium such as having: a cell wall, a cytoplasmic membrane, a nucleoid, cytoplasm, ribosomes, pili, and flagella^{11,12}. The overall shape of *L. acidophilus* is one that is rod shaped. This bacterium occurs in small chains and is usually 0.5 to 0.8 micrometre (1 mm=10⁻⁶ metre) across by 2 to 9 mm in length. It is also important to note that *L. acidophilus* is non-spore-forming.

Table 1: Fresh weight of curd bacteria in gm in presence of different types of sugars after 1 day by inoculation using spread plate method

S. No.	Types of sugar	Weight of curd bacteria in gm
1.	Nil sugar added (simple NA)	0.98 gm
2.	Sucrose	6.08 gm
3.	Fructose	3.28 gm
4.	Dextrose	2.17 gm
5.	Lactose	10.02 gm



Graph 1: Weight of curd bacteria in grams obtained by inocula containing different types of sugars

It was observed according to table 1 that the weight of curd bacteria was maximum in case of presence of lactose, followed by sucrose and fructose.

CONCLUSIONS AND FUTURE SCOPE

Curd bacterial rate of growth can be increased by supplementation of lactose

The present study revealed that curd bacteria can be increased in their biomass and rate of growth by addition of lactose in the inoculums followed by fructose. This can cause more formation of curd in less time. The process done in vitro can be scaled up in large bioreactors¹³.

Curd has probiotic microbes utilized as most beneficial health supplements

Probiotics are microorganisms that are believed to provide health benefits when consumed. The term probiotic is currently used to name ingested microorganisms associated with benefits for humans and animals. The term came into more common use after 1980. The introduction of the concept is generally attributed to Nobel recipient Élie Metchnikoff, who in 1907 suggested that "the dependence of the intestinal microbes on the food makes it possible to adopt measures to modify the flora in our bodies and to replace the harmful microbes by useful microbes". A significant expansion of the potential market for probiotics has led to higher requirements for scientific substantiation of putative benefits conferred by the microorganisms^{14,15,16}.

Commonly claimed benefits of probiotics include the decrease of potentially pathogenic gastrointestinal microorganisms, the reduction of gastrointestinal discomfort, the strengthening of the immune system, the improvement of the skin's function, the improvement of bowel regularity, the strengthening of the resistance to cedar pollen allergens, the decrease in body pathogens, the reduction of flatulence and

bloating, the protection of DNA, the protection of proteins and lipids from oxidative damage, and the maintaining of individual intestinal microbiota in subjects receiving antibiotic treatment^{17,18,19}.

Scientific evidence to date has been insufficient to substantiate any antidisease claims or health benefits from consuming probiotics^{20,21}.

REFERENCES

- [1] Alvarez-Olmos, M.I., Oberhelman, R.A.; Oberhelman (2001). "Probiotic agents and infectious diseases: a modern perspective on a traditional therapy". *Clin. Infect. Dis.* **32** (11): 1567–76.
- [2] Bâati, L. L.; Fabre-Gea, C.; Auriol, D.; Blanc, P. J. (2000). "Study of the cryotolerance of *Lactobacillus acidophilus*: Effect of culture and freezing conditions on the viability and cellular protein levels". *International Journal of Food Microbiology* **59** (3): 241–247.
- [3] de Roos N., Katan, M. (1 February 2000). "Effects of probiotic bacteria on diarrhea, lipid metabolism, and carcinogenesis: a review of papers published between 1988 and 1998". *Am J Clin Nutr* **71** (2): 405–11.
- [4] Fijan, Sabina (2014). "Microorganisms with Claimed Probiotic Properties: An Overview of Recent Literature". *International Journal of Environmental Research and Public Health* **11** (5): 4745–4767.
- [5] Fuller, R., ed. (1992) *Probiotics. The Scientific Basis*. Chapman & Hall, London, U.K
- [6] Gismondo MR, Drago L, Lombardi A; Drago; Lombardi (1999). "Review of probiotics available to modify gastrointestinal flora". *Int. J. Antimicrob. Agents* **12** (4): 287–92.
- [7] Glucose Tests.": The Test. American Association for Clinical Chemistry, n.d. Web. 17 Nov. 2015.
- [8] Hyvonen, L., & Koivistoinen, P (1982). "Fructose in Food Systems". In Birch, G.G. & Parker, K.J. *Nutritive Sweeteners*. London & New Jersey: Applied Science Publishers. pp. 133–144.
- [9] Kretchmer, N; Hollenbeck CB (1991). "Sugars and Sweeteners". CRC Press, Inc.
- [10] "Lactose Intolerance". National Digestive Diseases Information Clearinghouse (NDDIC). NIDDK. Retrieved 29 November 2011.
- [11] Lemieux, R. U.; Huber, G. (1953). "A chemical synthesis of sucrose". *J. Am. Chem. Soc.* **75** (16): 4118.
- [12] Ljungh A, Wadström T (2006). "Lactic acid bacteria as probiotics". *Curr Issues IntestMicrobiol* **7** (2): 73–89.
- [13] Mach T (November 2006). "Clinical usefulness of probiotics against chronic inflammatory bowel diseases". *Journal of Physiology and Pharmacology*. **57** Suppl 9: 23–33.

- [14] Makarova, K.; Slesarev, A.; Wolf, Y.; Sorokin, A.; Mirkin, B.; Koonin, E.; Pavlov, A.; Pavlova, N.; et al. (Oct 2006). "Comparative genomics of the lactic acid bacteria". *Proc Natl Acad Sci U S A* **103** (42): 15611–6.
- [15] Mintz, Sidney (1986). *Sweetness and Power: The Place of Sugar in Modern History*. Penguin.
- [16] O'Connell S, Walsh G (2006). "Physicochemical characteristics of commercial lactases relevant to their application in the alleviation of lactose intolerance". *Appl. Biochem. Biotechnol.* **134** (2): 179–91
- [17] Petrova, Mariya I.; Lievens, Elke; Malik, Shweta; Imholz, Nicole; Lebeer, Sarah (2015). "Lactobacillus species as biomarkers and agents that can promote various aspects of vaginal health". *Frontiers in Physiology* **6**.
- [18] Rettger, L.F., W.N. Levy, L. Weinstein, and J.E. Weiss. 1935. *Lactobacillus acidophilus and its therapeutic application*. Yale University Press, New Haven
- [19] Rijkers GT, de Vos WM, Brummer RJ, Morelli L, Corthier G, Marteau P; De Vos; Brummer; Morelli; Corthier; Marteau (2011). "Health benefits and health claims of probiotics: Bridging science and marketing". *British Journal of Nutrition* **106** (9): 1291–6.
- [20] Vesa TH, Marteau P, Korpela R (2000). "Lactose intolerance". *J Am Coll Nutr***19**: 165S–175S.
- [21] Wong SY, Hartel RW.: Crystallization in lactose refining-a review. *J Food Sci* 2014, 79(3):R257-72.[
- [22] Balamurugan R, Chandragunasekaran AS, Chellappan G, Rajaram K, Ramamoorthi G, Ramakrishna BS. Probiotic potential of lactic acid bacteria present in home made curd in southern India. *Indian J Med Res.* 2014 Sep;140(3):345-55. PMID: 25366201; PMCID: PMC4248380.
- [23] Suharman et. al 2021 *J. Phys.:* Conf. Ser. 1823 012038
- [24] Diaz, M., Del Rio, B., Ladero, V., Redruello, B., Fernández, M., Martín, M. C., et al. (2020). Histamine production in *Lactobacillus vaginalis* improves cell survival at low pH by counteracting the acidification of the cytosol. *Int. J. Food. Microbiol.* 321:108548. doi: 10.1016/j.ijfoodmicro.2020.108548
- [25] Zhang, Y., Vadlani, P., Kumar, A., Hardwidge, P., Govind, R., Tanaka, T., et al. (2016). Enhanced D-lactic acid production from renewable resources using engineered *Lactobacillus plantarum*. *Appl. Microbiol. Biot* 100, 279–288. doi: 10.1007/s00253-015-7016-0