Literature Review on Extrasolar Planets (Exoplanets)

Anusha S*

Amrita Vishwa Vidyapeetham Amritanagar PO, Ettimadai, Coimbatore

Abstract

The discovery of exoplanets since 1995 has revolutionized our understanding of planets and planetary systems. This paper reviews major detection techniques such as radial velocity, transit photometry, direct imaging, gravitational microlensing, astrometry, and pulsar timing, focusing on their strengths, limitations, and findings. It also highlights significant missions like Kepler, TESS, Gaia, and future contributions of the JWST. A synthesis of the literature reveals the critical role of interdisciplinary approaches and advancing technology in the ongoing search for habitable exoplanets and extraterrestrial life. This paper has also discussed about Exomoons and Exorings, which are less explored, but an advancing area in the exoplanetary science.




Figure 1: Comparison of different exoplanets with Earth

1. Introduction

Exoplanets, or extrasolar planets, orbit stars beyond our solar system. Since the first confirmed detection in 1992, over 5,000 exoplanets have been discovered. This review evaluates key detection techniques, significant findings, and the evolving understanding of planetary diversity and habitability.

2. Historical Background

Exoplanet science began in earnest with the discovery of pulsar planets in 1992 and the radial velocity detection of 51 Pegasi b in 1995. These milestones ushered in a new era of astronomy, enabling the systematic study of planetary systems.

3. Types of Exoplanets

- 1.) **Gas Giants:** Massive planets like Jupiter, with thick hydrogen-helium atmospheres.
- 2.) **Hot Jupiters:** Gas giants with close-in orbits, experiencing intense heat.
- 3.) **Super-Earths:** Planets with masses between Earth and Neptune, potentially rocky.
- 4.) **Mini-Neptunes:** Smaller than Neptune, with substantial atmospheres.
- 5.) **Terrestrial Planets:** Earth-like rocky planets, potentially habitable.
- 6.) **Ice Giants:** Neptune-like planets with icy cores beneath thick atmospheres.
- 7.) **Rogue Planets:** Free-floating planets untethered to stars.

4. Detection Techniques

4.1 Radial Velocity

Measures stellar wobble due to gravitational pull of orbiting planets. Effective for detecting large, close-in planets but limited for smaller or distant planets. Notable Discovery: 51 Pegasi b.

4.2 Transit Photometry

Detects brightness dips as planets transit their stars, allowing size, orbit, and atmospheric composition analysis. Requires specific orbital alignments. Notable Discovery: Kepler-186f.




Figure 2: Transit Photometry Detection Technique

4.3 Direct Imaging

Uses advanced optics to capture images of planets by blocking starlight. Effective for young, distant planets but technically challenging. Notable Discovery: HR 8799 planetary system.

4.4 Gravitational Microlensing

Relies on gravitational lensing effects to detect distant planets. Useful for wide orbits but limited by rare observation opportunities. Notable Discovery: OGLE-2005-BLG-390Lb.




Figure 3: Gravitation Microlensing Explanation

4.5 Astrometry

Tracks tiny positional shifts in stars caused by orbiting planets. Provides precise mass and orbit data but demands exceptional precision. Notable Discovery: HD 176051 b.

4.6 Pulsar Timing

Measures timing variations in pulsar signals caused by orbiting planets. Highly precise but restricted to pulsar systems. Notable Discovery: PSR B1257+12.




Figure 4: Pulsar Timing Detection Technique's Depiction

5. Exomoons

Exomoons are moons orbiting exoplanets, analogous to natural satellites in our Solar System like Earth's Moon or Jupiter's Europa. These could exist around planets in a wide range of environments, including gas giants, ice giants, and even terrestrial exoplanets.

Importance of Exomoons

- Habitability Potential: Large exomoons within the habitable zone of their star could retain atmospheres and liquid water, making them potential candidates for life.
- Insight into Planetary Formation: Studying exomoons helps understand the formation and evolution of planetary systems.
- Indicators of Planetary Properties: Detecting moons can reveal information about the planet's mass, orbit, and history.

Detection Techniques

- 1.) Transit Timing Variations (TTV) and Transit Duration Variations (TDV): The gravitational influence of a moon causes periodic variations in the timing and duration of the planet's transit.
- 2.) Direct Imaging: Advanced telescopes may be able to resolve large, distant exomoons directly.
- 3.) Microlensing: A moon-planet system can create unique microlensing light curves, although these events are rare.
- 4.) Spectroscopy: Spectral signatures during transit could reveal a moon's atmosphere, if it exists.

Challenges in Detection

Small size relative to the host planet makes exomoons faint and difficult to detect. Current detection limits make finding Earth-sized or smaller moons particularly challenging.

Notable Research and Candidates

Kepler-1625b-i: A potential exomoon around a Jupiter-sized exoplanet. This candidate is large (about Neptune-sized) and has sparked significant interest but remains unconfirmed.

MOA-2011-BLG-262: A microlensing event suggesting a moon orbiting a free-floating planet, although this too requires further confirmation.

Future Prospects

Advanced missions like the James Webb Space Telescope (JWST) and future observatories (e.g., ELT, Roman Space Telescope) will refine our ability to detect and study exomoons.

6. Exorings

Exorings are ring systems surrounding exoplanets, similar to Saturn's rings in our Solar System. They could also exist around brown dwarfs or even stars.

Importance of Exorings

- Planetary Evolution: Rings provide clues about the planet's formation, age, and environment.
- Habitability Influence: Rings can affect the stability of a planet's climate or impact the detectability of moons.

• Unique Observational Opportunities: Rings can enhance or obscure transit signals, offering new challenges and opportunities in exoplanet detection.

Detection Techniques

- 1. **Transit Light Curves:** Rings create distinctive light curves during transits, including "dips" and "shoulders" around the planet's main transit event.
- 2. **Polarimetry:** Ring systems can produce detectable polarization in the starlight scattered by the rings.
- 3. **Gravitational Microlensing:** As with exomoons, rings can create unique patterns in microlensing events.

Notable Research and Candidates

- **J1407b:** A giant exoplanet candidate with an enormous and complex ring system, estimated to be over 200 times the size of Saturn's rings.
- **Beta Pictoris:** Observations of debris disks in this system suggest potential ring systems forming around planets.

Challenges in Detection

- Rings are faint and require high-resolution data for unambiguous detection.
- Complex dynamics between rings, moons, and planets can make modeling difficult.
- Formation and Stability
- Rings are believed to form from debris left over after planet formation or from collisions.
- Tidal forces and gravitational interactions play a key role in their evolution and stability.

Future Prospects

Next-generation telescopes will enhance the resolution required to observe faint ring systems. Spectroscopic studies could identify the composition of ring material, such as ice or dust.

Exomoons and exorings are often interconnected, as moons can help stabilize rings, and rings can impact moon formation and dynamics. Studying these features together can reveal complex gravitational and orbital interactions.

7. Notable Missions And Surveys

- [1] Kepler: Identified over 2,600 exoplanets, emphasizing Earth-sized and habitable-zone candidates.
- [2] TESS: Continues Kepler's legacy, focusing on nearby stars. Key discovery: TOI-700d.
- [3] Gaia: Astrometric mission expected to uncover thousands of exoplanets.
- [4] JWST: Designed for detailed atmospheric studies of exoplanets.



Figure 5: Kepler Space Mission

8. Future Directions

Advancements in telescope technology, data analysis, and interdisciplinary styles are set to expand exoplanetary wisdom. A primary focus will be relating inhabitable exoplanets and implicit biosignatures.

Exomoons and exorings also represent promising areas of unborn disquisition, bettered discovery ways, similar as conveyance timing variations, high- resolution imaging, and advanced spectroscopic styles, will enhance our capability to identify and characterize exomoons. These moons could crop as crucial campaigners for habitability, particularly in cases where they circumvent gas titans within the inhabitable zone. Exorings, on the other hand, will give new perceptivity into the processes governing planetary conformation and elaboration. Detecting and studying these structures will exfoliate light on the dynamics of planetary systems and their commerce with girding debris.

With the development of coming - generation lookouts similar as the Roman Space Telescope, ELT, and ARIEL, the eventuality to descry lower, Earth - suchlike moons and faint exoring systems will significantly increase. These advancements would not

only consolidate our understanding of planetary systems but also help upgrade models of habitability, making the study of exomoons and exorings an essential part of unborn exoplanetary wisdom.

Conclusion

Exoplanet research has redefined planetary science, uncovering a diversity of worlds and hinting at the possibility of life beyond Earth. The study of exoplanets has been complemented by the emerging exploration of exomoons and exorings, which add further dimensions to our understanding of planetary systems. Exomoons, with their potential for habitability and unique dynamical interactions, and exorings, which provide insights into planetary formation and evolution, represent exciting frontiers in this field. With rapid technological progress, advanced detection methods, and collaborative scientific efforts, the next decades promise transformative discoveries, including the potential identification of habitable environments and the unraveling of complex planetary architectures.

References

Articles:

- [1] Beaulieu, J. P., et al. (2006). Discovery of a cool planet of 5.5 Earth masses through gravitational microlensing. Nature, 439(7075), 437-440.
- [2] Borucki, W. J., et al. (2014). Kepler-186f: A Potentially Habitable Planet in the Kepler-186 System. Science, 344(6181), 277-280.
- [3] Mayor, M., & Queloz, D. (1995). A Jupiter-mass companion to a solar-type star. Nature, 378(6555), 355-359.
- [4] Marois, C., et al. (2008). Direct Imaging of Multiple Planets Orbiting the Star HR 8799. Science, 322(5906), 1348-1352.
- [5] Muterspaugh, M. W., et al. (2010). The Phases Differential Astrometry Data Archive. The Astronomical Journal, 140(5), 1623-1637.
- [6] Wolszczan, A., & Frail, D. A. (1992). A planetary system around the millisecond pulsar PSR1257 + 12. Nature, 355(6356), 145-147.

Image Citations

• Fig 1: Comparison of different exoplanets with Earth

Source: https://www.futurelearn.com/courses/icy-moons-and-exoplanets

• Fig 2: Transit Photometry Detection Technique

 $Source: \underline{https://www.skyatnightmagazine.com/space-science/exoplanets-transit-\underline{method}}$

• Fig 3: Gravitation Microlensing Explanation

Source: https://www.eurekalert.org/news-releases/920957

• Fig 4: Pulsar Timing Detection Technique's Depiction

Source: https://www.youtube.com/watch?v="0mZblOlb2E

• Fig 5: Kepler Space Mission

Source: https://en.wikipedia.org/wiki/Kepler_space_telescope